Supplementary Data

Lipid Oxidation and Modification of Amyloid- β (A β) *in vitro* and *in vivo*

Grace Ellis^a, Erica Fang^a, Mukesh Maheshwari^a, Emily Roltsch^b, Leigh Holcomb^c, Danna Zimmer^b, Daniel Martinez^d and Ian V.J. Murray^{a,*}

^aNeuroscience and Experimental Therapeutics, College of Medicine, Texas A&M Health Science Center, College Station, TX, USA

^bVeterinary Pathobiology, Texas A&M University, College Station, TX, USA

^cCentral Texas Veterans Health Care System Neuropsychiatry Research Program, Temple, TX, USA

^d Pathology Core laboratory, Children's Hospital of Philadelphia, Philadelphia, PA, USA

Accepted 7 July 2010

^{*}Correspondence to: Dr. Ian V.J. Murray, 142E Reynolds Medical Building, Department of Neuroscience and Experimental Therapeutics, Texas A&M Health Science Center, College Station, TX 77843-1114, USA. Tel.: +1 979 458 4355; Fax: +1 979 845 0790; E-mail: IVMurray@medicine.tamhsc.edu.

G. Ellis et al. / Lipid Oxidation and $A\beta$



Supplemental Fig. 1. Antigen retrieval with 70% Formic acid abolishes HNE-adduct immunoreactivity. A) A β immunostaining in the hippocampus of saggital brain sections from a 12mo old PSAPP. B) Pretreatment of a near serial section with 70% formic acid markedly reduces the A β immunostaining. Magnification 10X and the scale bar represents 50 μ m.