

## Supplementary Data

---

# The Role of CNI-1493 in the Function of Primary Microglia with Respect to Amyloid- $\beta$

Jan-Philipp Bach<sup>a,1</sup>, David Mengel<sup>a,1</sup>, Tina Wahle<sup>b</sup>, Andreas Kautz<sup>a</sup>, Monika Balzer-Geldsetzer<sup>a</sup>, Yousef Al-Abed<sup>c</sup>, Richard Dodel<sup>a,\*</sup> and Michael Bacher<sup>a</sup>


<sup>a</sup>*Department of Neurology, Philipps-University Marburg, Marburg, Germany*

<sup>b</sup>*Department of Human Genetics, Laboratory of Neuronal Differentiation, Leuven, Belgium*

<sup>c</sup>*Laboratory of Medicinal Chemistry, The Feinstein Institute for Medical Research, Manhasset, NY, USA*

Handling Associate Editor: Brit Mollenhauer

Accepted 18 March 2011



Supplementary Figure 1. Cells were treated with LPS (1  $\mu$ g/ml) or 5  $\mu$ M CNI-1493 and the same concentration of LPS. In addition, untreated cells were used as negative control. IL-6 concentrations were measured using ELISA Assay. Addition of LPS led to a significant increase in IL-6 secretion, whereas CNI-1493 reversed this increase in cytokine secretion.

---

<sup>1</sup> Both authors contributed equally.

\*Correspondence to: Prof. Dr. Richard Dodel, Department of Neurology, Philipps-University Marburg, Baldingerstrasse, 35043 Marburg, Germany. Tel.: +49 6421-5866251; Fax: +49 6421 5865474; E-mail: dodel@med.uni-marburg.de.