

Supplementary Data

FOXP2, APOE, and PRNP: New Modulators in Primary Progressive Aphasia

Enrico Premi^{a,1}, Andrea Pilotto^{a,1}, Antonella Alberici^a, Alice Papetti^a, Silvana Archetti^b, Davide Seripa^c, Antonio Daniele^d, Carlo Masullo^d, Valentina Garibotto^e, Barbara Paghera^f, Federico Caobelli^f, Alessandro Padovani^a and Barbara Borroni^{a,*}

^aCentre for Ageing Brain and Neurodegenerative Disorders, Neurology Unit, University of Brescia, Brescia, Italy

^bLaboratories of Biotechnology, Brescia Hospital, Brescia, Italy


^cDepartment of Medical Sciences, Istituto Di Ricovero e Cura a Carattere Scientifico, Casa Sollievo della Sofferenza, San Giovanni Rotondo (FG), Italy

^dNeuropsychology Unit, Catholic University, Rome, Italy

^eDepartment of Medical Imaging, Geneva University Hospital, Geneva, Switzerland

^fNuclear Medicine Unit, University of Brescia, Brescia, Italy

Accepted 4 November 2011



Supplementary Figure 1. Reduction in regional cerebral perfusion in Primary progressive aphasia patients compared to a group of healthy subjects, showing a prevalent left frontotemporal hypoperfusion ($n = 14$, % Female = 36% (5), mean age = 58.9 years, SD = 11.8 years). The threshold was set at $p < 0.05$ corrected for multiple comparisons (FDR). L = left.

¹These authors equally contributed to this work.

*Correspondence to: Barbara Borroni, MD, Neurology Unit, University of Brescia, Piazza Spedali Civili 1, 25125 Brescia, Italy.
E-mail: bborroni@inwind.it.