

Supplementary Data

TOMM40, APOE, and APOC1 in Primary Progressive Aphasia and Frontotemporal Dementia

Davide Seripa^{a,*}, Alessandra Bizzarro^b, Andrea Pilotto^c, Orazio Palmieri^d, Francesco Panza^a, Grazia D'Onofrio^a, Carolina Gravina^a, Silvana Archetti^e, Antonio Daniele^b, Barbara Borroni^c, Alessandro Padovani^c and Carlo Masullo^b

^a*Gerontology and Geriatrics Research Laboratory, I.R.C.C.S. "Casa Sollievo della Sofferenza", San Giovanni Rotondo (FG), Italy*

^b*U.O.C. of Neurology, "Agostino Gemelli" General Hospital, Catholic University School of Medicine, Rome, Italy*

^c*Centre for Ageing Brain and Neurodegenerative Disorders, Neurology Unit, University of Brescia, Brescia, Italy*


^d*Gastroenterology Research Laboratory, I.R.C.C.S. "Casa Sollievo della Sofferenza", San Giovanni Rotondo (FG), Italy*

^e*Laboratories of Biotechnology, Azienda Ospedaliera "Spedali Civili", Brescia, Italy*


Handling Associate Editor: Daniela Galimberti

Accepted 3 May 2012

*Correspondence to: Dr. D. Seripa, Gerontology and Geriatrics Research Laboratory, I.R.C.C.S. "Casa Sollievo della Sofferenza", Viale Cappuccini 1, 71013 San Giovanni Rotondo (FG), Italy. Tel.: +39 0882 416260; Fax: +39 0882 416 264; E-mail: dseripa@operapadrepio.it.



Supplementary Figure 1. Linkage disequilibrium blocks at 19q13-q13.2. The tag SNPs ($D' > 0.8$) chosen for the analysis are circled in red.



Supplementary Figure 2. Schematic representation of r^2 linkage disequilibrium coefficient at the 26.9-kb block spanning the 19q13-q13.2 chromosomal region in frontotemporal lobar degeneration (FTLD) (A) and in FTLD-subtypes primary progressive aphasia (PPA) and behavioral variant frontotemporal dementia (bvFTD) (B).