

Supplementary Data

Natural Cannabinoids Improve Dopamine Neurotransmission and Tau and Amyloid Pathology in a Mouse Model of Tauopathy

Maria J. Casarejos^{a,e,1}, Juan Perucho^{a,e,1}, Ana Gomez^{a,e}, Maria P. Muñoz^{a,e}, Marian Fernandez-Estevez^{a,e}, Onintza Sagredo^{c,e}, Javier Fernandez Ruiz^{c,e}, Manuel Guzman^{d,e}, Justo Garcia de Yébenes^{b,e} and Maria A. Mena^{a,e,*}

^a*Departments of Neurobiology, Ramon y Cajal University Hospital, Madrid, Spain*

^b*Departments of Neurology, Ramon y Cajal University Hospital, Madrid, Spain*

^c*Department of Biochemistry and Molecular Biology, School of Medicine, UCM, Madrid, Spain*


^d*Department of Biochemistry and Molecular Biology, School of Biology, UCM, Madrid, Spain*

^e*CIBERNED, Spain*


Accepted 4 February 2013

¹These authors contributed equally to this work.

*Correspondence to: Dr. M.A. Mena, Department of Neurobiology, Hospital Ramón y Cajal, Ctra. de Colmenar, Km. 9, Madrid 28034, Spain. Tel.: +34 91 336 83 84; Fax: +34 91 336 90 16; E-mail: maria.a.mena@hrc.es.



Supplementary Figure 1. Breeding mice schema and PCR genotyping to achieve the parkin-null, human tau overexpressing ($PK^{-/-}/Tau^{VLW}$) mice (% expected/% found).



Supplementary Figure 2. Tau and amyloid-β ($A\beta$) pattern in $PK^{-/-}/Tau^{VLW}$ and wild-type (WT) age-matched mice and biochemical characterization of soluble $A\beta$ oligomers (ADDLs) followed by western blot (WB) with 6E10.